

WINEMAKING GUIDELINES

RED COMPROMISED FRUITS: POWDERY MILDEW

Powdery mildew is a fungal disease that affects grapevines, caused by the fungus *Erysiphe necator*. If not controlled, powdery mildew can cause crop losses and have negative impacts on wine quality. The presence of powdery mildew can also increase the risk of Botrytis infection. The contamination of grapes with powdery mildew can have many challenges for winemakers:

- Sensitivity to oxidation
- Stuck fermentations due to nutrient deficiencies, toxins accumulation and microbial contamination
- Fragile grapes, sensitive to secondary contaminations
- Off-aromas and flavors such as mushroom, earthy, wet fur and cooked tomato characters
- High protein content due to pathogenesis-related protein expression
- Increases 'oiliness' and viscosity sensation in the palate
- High pH, high TA
- High phenolic content and bitterness
- Color instabilities.

KEY WINEMAKING STEPS WHEN DEALING WITH POWDERY MILDEW AFFECTED GRAPES:

- Hand harvest and sort contaminated grapes in the vineyard and at the winery.
- Control any spoilage microbes as early as possible with SO2 and bio-protection with Excellence B-Nature
- Use adequate antioxidant protection to limit browning, color loss and aroma oxidation such as Pro Tanin R
- Reduce skin contact to limit extraction of off-flavors, no cold soak, avoid extended maceration, use short pressing cycles and separate press fractions (first 10 gallons).
- Fining and clarification of juice to eliminate off-aromas and 'clean' the must for good fermentation conditions. We recommend using Polymix Natur' for an efficient fining during a delestage (reds) done in the early stage of fermentation.
- Supplement must with vitamins, amino acids and ammonia to ensure complete healthy fermentation
- Boost fresh fruit aromatic expression by using specific yeast nutrient such as **OptiEsters**
- Improve color intensity and stability by promoting condensation and co-pigmentation reactions during fermentation with <u>Softan</u> Vinification
- Balance phenolic compounds early in fermentation with Natur'Soft
- Consider fining, mannoproteins or tannins additions to balance mouthfeel and remove aggressive phenolics.
- During ageing, it is essential to balance the redox potential of the wine and protect from microbial development to prevent premature ageing and spoilage.



WINEMAKING GUIDELINES

FOCUS PRODUCTS

Excellence B-Nature — non-Saccharomyces yeast, pure *Metschnikovia pulcherrima*, non-fermentative. It inhibits the development of spoilage microbes such as other non-Saccharomyces, and bacteria on grapes and juice. Excellence® B-Nature® is an organic anti-microbial solution, used as alternative to SO2 on grapes. It protects grapes/juice from microbial contamination during transport and processing, does not inhibit Saccharomyces cerevisiae, and reduces SO2 combining compounds production, thus increasing SO2 efficiency. Excellence B-nature can be added directly to grapes, without rehydration. Simply sprinkle the yeast on the top of the grapes at picking.

<u>Pro Tanin R</u> – Pro-anthocyanidin tannin, use as sacrificial tannin. Developed for application on red grapes, to scavenge oxygen radicals, inhibit oxidative enzymes such as laccase and PPO and eliminates reactive proteins, thus protecting grape polyphenols. It is instantaneously soluble, simply sprinkle it on the top of the grapes at harvest.

<u>Polymix Natur'</u> – PVPP, Yeast extracts, Bentonite. Vegan, allergen free fining agent focused on removing oxidized and easily oxidable phenolic compounds. Polymix Natur' treats and prevents oxidation, improves oxidative stability, wine expression and elongates wine shelf life.

<u>Oenozym Clar</u> – Pectinase with strong de-pectinization activity for a quick and effective clarification. Purified from side activities such as cinnamyl-esterase and anthocyanases. Resistant to low (5°C) and high (68°C) temperatures.

<u>OptiEsters</u> - Yeast nutrient composed of amino acids and ergosterols selected to increase the production of esters, acetates and ethylesters, thus increase red fruits, floral, berries notes. It increases aromatic complexity, freshness and intensity. Interesting tools to boost freshness and complexity of a wine, as well as compensate for off-aromas.

<u>Softan Vinification</u> – catechins tannins bounded to plant polysaccharides. Added during fermentation, Softan Vinification has a strong ability to stabilize color and protect it from loss during fermentation. It is a gentle tannin that contributes to mouthfeel and build up mid-palate.

<u>Natur'Soft</u> - preparation of specific yeasts hulls, selected for their high content of polysaccharides. It is strongly effective in color stabilization, as well as filling mid palate and improving mouthfeel. It increases wine complexity, reduces tannins perception, and stabilize color.

<u>Vinitan Advance</u> – pure grape tannin with low phenol content. It improves wine structure while respecting its finesse and balance. Excellent for color stabilization during ageing, it also boosts the fruits aromas and maintain a clean and balance wine profile during ageing.

<u>KillBrett</u> – pure chitosan, wide spectrum anti-microbial agent. KillBrett eliminates and inhibits Brettanomyces, Lactic Acid Bacteria and Acetic Acid Bacteria. It can be used during the entire process of winemaking, we recommend using it as preventive, post MLF, at 4 g/hL

<u>Aroma Protect</u> - inactivated yeasts, naturally rich in glutathione, a natural antioxidant, sulfurous tripeptide with great reductive power. It gives immediate protection against oxidative mechanisms, releasing glutathione (GSH) into the wine, thus slowing down oxidation.

<u>Tan&Sense Volume</u> – pure untoasted oak tannins, with high capacity to scavenge oxygen radicals, buffer redox potential and maintain wine freshness. It is a gentle tannin, increasing sweetness and roundness perception.



WINEMAKING GUIDELINES

RED GRAPES COMPROMISED FRUIT WITH POWDERY MILDEW - WINEMAKING GUIDELINES

HARVEST AND GRAPE TRANSPORT	Rigorously sort, in the vineyard or on processing in the winery. Protect from oxidation and microbial contamination as soon as possible when picking to limit browning. - 40-60 ppm of SO ₂ - Add 180-220g/ton of Pro Tanin R, sprinkled on the grapes at harvest to protect from oxidation and prevent the loss of phenolic compounds. - Excellence B-Nature at 50 g/ton, sprinkle directly on grapes, as soon as possible after picking to prevent any microbial contamination and spoilage. In case of high contamination levels, thermovinification can be an option, please see our specific protocol.
MACERATION	Limit the use of extraction enzymes. Avoid cold soaking and extended macerations.
ALCOHOLIC FERMENTATION	Powdery Mildew contamination leads to nitrogen deficiencies which needs to be compensated to prevent stuck fermentation. It is essential rehydrate yeast with OenoStim at 30 g/hL to reinforce yeast activity, increase aromatic production and ensure yeast health and resistance along the fermentation. Yeast: Excellence DS at 20 g/hL to produce fresh, fruity, spicy and elegant profile with smooth structure OptiEsters at 15 g/hL to promote the production of ethylesters and enhance fresh, fruity and floral characters. 1 DAY AFTER INOCULATION At day 1, fine and clarify the juice to remove any off-aromas, undesirable phenolic compounds, signs of oxidation and toxins that will inhibit fermentation. Proceed to a delestage (drain all must) protected from oxygen. Add Polymix Natur' at 40-60 g/hL into the drained juice. Rack the 'cleaned' juice back on the top the grapes. Ensure good yeast nutrition and limit off-flavors production with Optiflore O® at 40 g/hL (complete organic nutrient based on inactivated yeast). Optiflore O also helps detoxifying the must for better fermentations. 120 g/ton of Softan Vinification to encourage the stabilization of anthocyanins via co-pigmentation and condensation, protect anthocyanins and fill mid palate. AT 18 BRIX Add 20-30 g/hL of OptiFerm (ammonium salts and vitamin B1) at 1/3 of fermentation. 150 g/ton of Natur'Soft to stabilize color, fill mid palate and increase wine length and volume.
PRESSING	Press early, when phenolic compounds extraction is sufficient (decide with tasting). - Add <u>Oenozym Clar</u> at 4 ml/hL to improve settling and clarification. Rack gross lees. - Add <u>Vinitan Advance</u> at 5 g/hL to improve wine phenolic balance, increase antioxidant resistance and stabilize color.
MLF	Add <u>Oeno1</u> at 1g/hL once AF is completed
AGEING	 Once AF and MLF completed: rack off gross lees after fermentation using inert gas. Add SO₂ 3-4 g/hL + <u>KillBrett</u> at 4 g/hL to stabilize wine from oxidation and microbial contamination Rebalance the redox potential and natural oxidation resistance of the wine by using <u>Aroma Protect</u> at 20 g/hL. It will maintain wine freshness, protect from oxidation, lower redox potential and limit SO₂ loss during ageing. Use <u>Tan&Sense Volume</u> at 1 g/hL (pure untoasted oak tannins) every racking to protect from oxidation, regulate redox potential and scavenge oxygen radicals. It will also give some roundness to wine.